

Special Local Need Registration

SUPPLEMENTAL LABELING FOR DISTRIBUTION AND USE ONLY IN MONTANA

Admire™ 2 Flowable

Insecticide

For control of certain insects infesting potato.

ACTIVE INGREDIENT:

Imidacloprid, 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine 21.4%

INERT INGREDIENTS 78.6%
100.0%

SHAKE WELL BEFORE USING

Contains 2 pounds of imidacloprid per gallon

EPA Reg. No. 3125-422

STOP - Read The Label Before Use
KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through skin.
Avoid contact with skin, eyes, or clothing.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Water-proof gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/ maintaining personal protective equipment, PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering controls statements:

- When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

STATEMENTS OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center. Drink one or two glasses of water and induce vomiting by touching back of throat with finger, or, if available, by administering syrup of ipecac. If syrup of ipecac is available, administer 1 tablespoonful (15 mL) of syrup of ipecac followed by 1 to 2 glasses of water. If vomiting does not occur within 20 minutes, repeat the dose once. Do not

induce vomiting or give anything by mouth to an unconscious person. **If on skin:** Wash thoroughly with soap and water. Get medical attention if irritation occurs. **If in eyes:** Hold eyelids open and flush with plenty of water.

To Physician: No specific antidote is available. Treat the patient symptomatically.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. This product is highly toxic to aquatic invertebrates.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

IMPORTANT: Read these entire DIRECTIONS FOR USE, AND CONDITIONS OF SALE before using ADMIRE 2 Flowable Systemic Insecticide.

CONDITIONS OF SALE: THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE APPROPRIATE FOR THE CORRECT USE OF THIS PRODUCT. THIS PRODUCT HAS BEEN TESTED UNDER DIFFERENT ENVIRONMENTAL CONDITIONS BOTH INDOORS AND OUTDOORS UNDER CONDITIONS SIMILAR TO THOSE THAT ARE ORDINARY AND CUSTOMARY WHERE THE PRODUCT IS TO BE USED. INSUFFICIENT CONTROL OF PESTS OR PLANT INJURY MAY RESULT FROM THE OCCURRENCE OF EXTRAORDINARY OR UNUSUAL CONDITIONS, OR FROM FAILURE TO FOLLOW LABEL DIRECTIONS. IN ADDITION, FAILURE TO FOLLOW LABEL DIRECTIONS MAY CAUSE INJURY TO ANIMALS, MAN, AND DAMAGE

TO THE ENVIRONMENT. BAYER OFFERS, AND THE BUYER ACCEPTS AND USES, THIS PRODUCT SUBJECT TO THE CONDITIONS THAT EXTRAORDINARY OR UNUSUAL ENVIRONMENTAL CONDITIONS, OR FAILURE TO FOLLOW LABEL DIRECTIONS ARE BEYOND THE CONTROL OF BAYER AND ARE, THEREFORE, THE RESPONSIBILITY OF THE BUYER.

Do not apply this product through any type of irrigation system unless it is specified in the "Recommended Application" section.

RESISTANCE: Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product should conform to resistance management strategies established for the use area. Consult your agricultural advisor for resistance management strategies and recommended pest management practices for your area.

ENDANGERED SPECIES NOTICE: Under the Endangered Species Act, it is a Federal Offense to use any pesticide in a manner that results in the death of a member of an endangered species. Consult your local county bulletin, County Extension Agent, or Pesticide State Lead Agency for information concerning endangered species in your area.

ROTATIONAL CROPS

Treated areas may be replanted with any crop specified on an imidacloprid label, or any crop for which a tolerance exists for the active ingredient, as soon as practical following the last application.

For crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12-month plant-back interval should be observed.

ROTATIONAL PLANT-BACK INTERVALS¹

Immediate Plant-Back		
Barley	Cauliflower	Pepinos
Broccoli	Collards	Pepper
Broccoli, Chinese	Cotton	Potatoes
Broccoli raab	Eggplant	Rape greens
Brussels Sprouts	Ground cherry	Sorghum
Cabbage, Chinese mustard	Kale	Sugarbeets
Cabbage, Chinese	Kohlrabi	Tomatillo
Cabbage	Lettuce	Tomato
Canola	Mustard greens	Wheat

30-Day Plant-Back
Cereals (Including buckwheat, corn, millet, oats, popcorn, rice, rye, and triticale)
Legumes (including soybeans, beans and peas)
Safflower

12-Month Plant-Back
All Other Crops

¹ Cover crops for soil building or erosion control may be planted any time, but do not graze or harvest for food or feed.

Special Local Need

Notification of the crop rotational restrictions must be conveyed to the grower through the distribution of this label on all commercially treated seed.

NOTE: Federal regulations have established official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates. Do not over treat ADMIRE-treated seed-pieces with GAUCHO, ADMIRE, or PROVADO

Labels for commercially treated seed should include the following addition to the Environmental Hazards statements:

Dispose of all excess treated seed-pieces and seed packaging by burial away from bodies of water. Cover or incorporate spilled treated seed-pieces.

RECOMMENDED APPLICATIONS			
CROP	PEST	RATE PER APPLICATION	REMARKS
Potato	Aphids Colorado potato beetle Flea beetles Potato leafhopper Psyllids Wireworms (seed-piece protection only)	Seed-Piece Application 0.4 to 0.8 fl oz per 100 lbs of seed-pieces (8 to 16 fl oz per acre)*	Apply specified dosage as a non-diluted or diluted spray onto seed-pieces using a shielded spray system. For dilute applications, do not dilute with any more than 3 parts water to 1 part ADMIRE 2 F. Agitate or stir spray solution as needed. Immediately after application, apply a fungicide that contains an inert absorbent ingredient (i.e. bark) or apply an inert absorbent ingredient to sprayed seed-pieces. Apply only in areas with adequate ventilation or in areas that are equipped to remove spray mist or dust. Plant seed-pieces as soon as possible after cutting and treating in accordance with the recommendations of your local extension potato specialist. Avoid prolonged exposure of ADMIRE-treated seed-pieces to sunlight. Do not use treated seed-pieces for food, feed, or fodder. The length of control will vary depending on the rate used. Consult your local Bayer or Gustafson representative or dealer for information relevant to your area. Note: Do not apply any subsequent application of imidacloprid (i.e. GAUCHO dust, ADMIRE-in-furrow, or PROVADO) following an ADMIRE seed-piece treatment

Exposed treated seed-pieces may be hazardous to birds.

* Based on a seeding rate of 2000 lb/acre.

This labeling must be in the possession of the user at the time of pesticide application.

EPA SLN No. MT-990001

Bayer Corporation
Crop Protection Products
Box 4913
Kansas City, MO 64120-0013

MTADSL99.028

IMPORTANT

Before using this product, read and carefully observe directions, cautionary statements and other information appearing on the product packaging label. This product is sold subject to the Conditions of Sale set forth on the container label.

